FINAL TECHNICAL REPORT ONR GRANT #N00014-91-J-1242

John A. Barth Buoyancy-Driven Coastal Currents

The stability of a coastal upwelling jet and front was investigated using the primitive equations applied to a continuously stratified flow in geostrophic balance (Barth, 1994). A linear stability analysis explained the growth of two modes of instability with distinctly different horizontal scales. A long-wavelength mode (fastest-growing wavelength of 0(100 km)) was a modified version of a traditional baroclinic instability. A second, rapidly growing frontal instability was also found and had the following properties: fastest growth at short wavelengths (0(20 km)); e-folded in less than 1.5 days; propagated rapidly in the direction of the mean flow. The frontal mode was trapped to the surface front and its influence was confined to the upper water column ($z \lesssim 70 \text{ m}$). A significant subsurface vertical velocity maximum (20 m d⁻¹ at 30 m) was associated with the frontal instability. The linear stability predictions were verified by and compared with results from a time-dependent, three-dimensional, nonlinear ocean circulation model.

The linear stability analysis was also applied to observed sections of salinity and velocity from the buoyancy-driven Alaska Coastal Current (Barth, 1994b,c). Again, two modes of mesoscale variability were found. One unstable mode represented a modified form of traditional baroclinic instability with fastest-growth (e-folding in 1.8 days) occurring at 63 km with a slow phase speed (0.09 m s⁻¹) in the direction of the mean flow. A second unstable ageostrophic, frontal instability mode had a fastest-growing wavelength of 40 km that e-folded in 2.3 days and propagated rapidly (0.57 m s⁻¹) in the direction of the mean flow. A three-dimensional, time-dependent, fine-resolution numerical ocean circulation model was used to study the nonlinear evolution of the unstable buoyancy-driven coastal jet. Early in the simulation, short-wavelength frontal instabilities grew and propagated along the jet. Eventually, the frontal instabilities were frictionally damped and the long-wavelength baroclinic instability became dominant. The wavelength (65 km) and phase speed (0.09 m s⁻¹) were in excellent agreement with linear predictions. The baroclinic instability amplified into a series of backward-breaking waves whose crests eventually pinched off to form fresh, anticyclones while saline cyclones remained in the troughs. The scales of these eddies (diameters of 40 km) were consistent with those evident in maps of dynamic height from the Alaska Coastal Current. Vigorous vertical velocities, as large as 25 m day⁻¹, associated with the unstable meandering coastal current were noted to be potentially important for the vertical flux of material into the euphotic zone in the Alaska Coastal Current.

Density and velocity data from a high-resolution, upper-ocean survey in the California Current using SeaSoar and a shipboard acoustic Doppler current profiler were used to diagnose vertical velocity via the quasigeostrophic omega equation (Shearman et al., 1998). The diagnosed vertical velocity field showed velocities of 30-40 m d⁻¹. The lateral distribution of relative vorticity and vertical velocity was characterized by patches with length scales of 20-30 km. Geostrophic streamline analysis of vertical velocity indicated that water parcels made net vertical excursions of 20-30 m over 2-3 days, resulting in net vertical velocities of

19980630 018

COLLEGE OF OCEANIC & ATMOSPHERIC SCIENCES

June 22, 1998



Oregon
State
University

Oceanography Adm Bldg 104 Corvallis, Oregon 97331.5503 Dr. Thomas H. Kinder Office of Naval Research Code 321CD, Room 428 800 North Quincy Street Arlington, VA 22217-5660

Dear Dr. Kinder,

In order to complete my ONR grant entitled "Buoyancy-Driven Coastal Currents", I am sending three copies of the *Final Technical Report* to you with copies distributed as indicated below, along with a completed *Report Documentation Page (SF 298)*.

Sincerely,

Dr. Jack A. Barth Associate Professor

Enclosures

Defense Technical Information Center (2) 8725 John J. Kingman Rd, Suite 0944 Ft Belvoir, VA 22060-6218

Administrative Contracting Officer (1) Office of Naval Research Seattle Regional Office 1107 NE 45th St, Suite 350 Seattle, WA 98105-4631

Director, Naval Research Laboratory (1) Attn: Code 2627 Washington, DC 20375

Telephone 541·737·1607

Fax 541.737.2064

10-15 m d⁻¹. Water parcels moving along geostrophic streamlines experienced maximum vertical velocities in regions of maximum alongstream change in relative vorticity, an indication of potential vorticity conservation.

LIST OF PUBLICATIONS

- Barth, J. A., 1994a. Short-wavelength instabilities on coastal jets and fronts. J. Geophys. Res., 99, 16095-16115.
- Pierce, S. D., J. A. Barth and R. L. Smith, 1998. Improving ADCP accuracy with WADGPS and adaptive smoothing of ship velocity. *J. Atmos. Oceanic Technol.*, submitted.
- Pierce, S. D., R. L. Smith, P. M. Kosro, J. A. Barth and C. D. Wilson, 1998. Continuity of the poleward undercurrent along the eastern boundary of the mid-latitude north Pacific. Deep-Sea Res., submitted.
- Shearman, R. K., J. A. Barth and P. M. Kosro, 1998. Diagnosis of the three-dimensional circulation associated with mesoscale motion in the California Current. *J. Phys. Oceanogr.*, in press.

Abstracts:

- Barth, J. A., 1992. Ageostrophic, short-wavelength instabilities on coastal jets and fronts. Eos, Transactions, American Geophysical Union, 72(51), 50.
- Barth, J. A., 1994b. Mesoscale variability generated by an unstable buoyancy-driven coastal current with application to the Alaska Coastal Current. Eos, Transactions, American Geophysical Union, 75(3), 133.
- Barth, J. A., 1994c. Mesoscale variability generated by an unstable buoyancy-driven coastal current with application to the Alaska Coastal Current. 7th International Biennial Conference on Physics of Estuaries and Coastal Seas, 28-30 November, Woods Hole, Ma.
- Pierce, S. D., J. A. Barth and P. M. Kosro, 1996. Applications of inexpensive Wide Area Differential GPS navigation. *Eos, Transactions, American Geophysical Union*, **77(3)**, Ocean Sciences Meet. Suppl., OS100.
- Pierce, S. D., R. L. Smith and P. M. Kosro, 1996. Observations of the poleward undercurrent along the eastern boundary of the mid-latitude Pacific. *Eos, Transactions, American Geophysical Union*, 77(46), Fall Meet. Suppl., F345.
- Shearman, R. K., J. A. Barth and P. M. Kosro, 1996. Diagnosis of three-dimensional circulation: Results from the Eastern Boundary Current program. *Eos, Transactions, American Geophysical Union*, 77(3), Ocean Sciences Meet. Suppl., OS131.
- Smith, R. L., S. D. Pierce, J. A. Barth, P. M. Kosro and C. D. Wilson, 1996. observations of the poleward undercurrent along the continental margin off California, Oregon and Washington. *Eos, Transactions, American Geophysical Union*, 77(3), Ocean Sciences Meet. Suppl., OS131.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson David Highest State 1204, Adjusting National Applications (Application Visit 2202-4302, and to the Office of Management and Budget Paperwork Reduction Project (0704-01188) Washington DC 20503

| Davis Highway, Suite 1204, Arlington, VA 22202-4302, | | | | |
|---|-----------------------------|----------------------------|--|--|
| 1. AGENCY USE ONLY (Leave blank) | 2. REPORT DATE 23 June 1998 | 3. REPORT TYPE AN Final | ND DATES COVERED | |
| 4. TITLE AND SUBTITLE | | | 5. FUNDING NUMBERS | |
| Buoyancy-Driven Coastal Currents | | | N00014-91-J-1242 | |
| 6. AUTHOR(S) | | | · | |
| John A. Barth | | | | |
| | | | · | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) | | | 8. PERFORMING ORGANIZATION | |
| Oregon State University | | | REPORT NUMBER | |
| Corvallis, OR 97331-550 |)3 | | | |
| | | | N00460 | |
| | | | | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) | | | 10. SPONSORING/MONITORING AGENCY REPORT NUMBER | |
| Office of Naval Research Ballston Centre Tower Or | | | N00014-91-J-1242 | |
| 800 North Quincy Street | | | N00014 91 0 1212 | |
| Arlington, VA 22217-566 | 50 | | | |
| | | | | |
| 11. SUPPLEMENTARY NOTES | | | | |
| | | | | |
| | | | | |
| 12a. DISTRIBUTION / AVAILABILITY STAT | EMENT | • | 12b. DISTRIBUTION CODE | |
| | | | | |
| Unlimited Public Access | | · | | |
| | | | | |
| | | | , | |
| 13. ABSTRACT (Maximum 200 words) | | | | |
| The stability of a coastal | upwelling jet and | front was inves | stigated using the primi- | |
| tive equations applied to a continuously stratified flow in geostrophic balance. A | | | | |
| linear stability analysis explained the growth of two modes of instability with dist- | | | | |
| inctly different horizontal scales. A long-wavelength mode was a modified version of | | | | |
| a traditional baroclinic instability. A second, rapidly growing frontal instability | | | | |
| was also found. The linear stability analysis was also applied to observed sections of | | | | |
| salinity and velocity from the buoyancy-driven Alaska Coastal Current and, again, two modes of mesoscale variability were found. A three-dimensional, time-dependent, fine- | | | | |
| resolution numerical ocean circulation model was used to study the evolution of the | | | | |
| unstable buoyancy-driven coastal jet. Early in the simulation, short-wavelength | | | | |
| frontal instabilities grew and propagated along the jet. Eventually, the frontal | | | | |
| instabilities were frictionally damped and the long-wavelength baroclinic instability | | | | |
| became dominant. The baroclinic instability amplified into a series of backward- | | | | |
| breaking waves whose crests eventually pinched off to form fresh, anti-cyclones while | | | | |
| saline cyclones remained in the troughs. The scales of these eddies were consisten | | | | |
| with those evident in maps of dynamic height from the Alaska Coastal Current. | | | | |

17. SECURITY CLASSIFICATION OF REPORT

unclassified

18. SECURITY CLASSIFICATION,
OF THIS PAGE
unclassified

19. SECURITY CLASSIFICATION OF ABSTRACT unclassified

20. LIMITATION OF ABSTRACT

14. SUBJECT TERMS

15. NUMBER OF PAGES

16.-PRICE CODE